

AN ALGORITHM FOR CONSTRUCTING CERTAIN DIFFERENTIAL OPERATORS IN POSITIVE CHARACTERISTIC

ALBERTO F. BOIX*, ALESSANDRO DE STEFANI†, AND DAVIDE VANZO

ABSTRACT. Given a non-zero polynomial f in a polynomial ring R with coefficients in a finite field of prime characteristic p , we present an algorithm to compute a differential operator δ which raises $1/f$ to its p th power. For some specific families of polynomials, we also study the level of such a differential operator δ , i.e., the least integer e such that δ is R^{p^e} -linear. In particular, we obtain a characterization of supersingular elliptic curves in terms of the level of the associated differential operator.

1. INTRODUCTION

Let $R = k[x_1, \dots, x_d]$ be the polynomial ring over a field k , and let \mathcal{D}_R be the ring of k -linear differential operators on R . For every non-zero $f \in R$, the natural action of \mathcal{D}_R on R extends uniquely to an action on R_f . In characteristic 0, it has been proven by Bernstein in the polynomial ring case (cf. [Ber72, Corollary 1.4]) that R_f has finite length as a \mathcal{D}_R -module. The minimal m such that $R_f = \mathcal{D}_R \cdot \frac{1}{f^m}$ is related to Bernstein-Sato polynomials (cf. [ILL⁺07, Theorem 23.7, Definition 23.8, and Corollary 23.9]), and there are examples in which $m > 1$ (e.g. [ILL⁺07, Example 23.13]). Remarkably, in positive characteristic, not only is R_f finitely generated as a \mathcal{D}_R -module [Bøg95, Proposition 3.3], but it is generated by $\frac{1}{f}$ (cf. [ÀMBL05, Theorem 3.7 and Corollary 3.8]). This is shown by proving the existence of a differential operator $\delta \in \mathcal{D}_R$ such that $\delta(1/f) = 1/f^p$, i.e., a differential operator that acts as the Frobenius homomorphism on $1/f$. The main result of this paper exhibits an algorithm that, given $f \in R$, produces a differential operator $\delta \in \mathcal{D}_R$ such that $\delta(1/f) = 1/f^p$. We will call such a δ a *differential operator associated with f* . Our method is described in full details in Section 3. Moreover, this procedure has been implemented using the computer algebra system Macaulay2.

Assume that $\text{char}(k) = p > 0$ and that $[k : k^p] < \infty$. For $e \geq 1$ let R^{p^e} be the subring of R consisting of all p^e -th powers of elements in R , which can also be viewed as the image of the e -th iteration of the Frobenius endomorphism $F : R \rightarrow R$. We set $R^{p^0} := R$. It is shown in [Yek92, 1.4.9] that \mathcal{D}_R is equal to the increasing union $\bigcup_{e \geq 0} \text{End}_{R^{p^e}}(R)$. Therefore, given $\delta \in \mathcal{D}_R$, there exists $e \geq 0$ such that $\delta \in \text{End}_{R^{p^e}}(R)$ but $\delta \notin \text{End}_{R^{p^{e'}}}(R)$ for any $e' < e$. Given a non-zero polynomial $f \in R$, we have seen above that there exists $\delta \in \mathcal{D}_R$ that is associated with f . We say that f has level e if such δ is R^{p^e} -linear, and there is no $R^{p^{e'}}$ -linear differential operator δ' , with $e' < e$, that is associated with f .

In Section 4, we study the case when f is a monomial; indeed, in Theorem 4.2 we determine the level of f , and we give an explicit description of the differential operator δ associated with f . We also describe explicitly $I_e(f^{p^e-1})$, the ideal of p^e -th roots of f^{p^e-1} , where e is the level of f . The ideal $I_e(f^{p^e-1})$ can be defined as the unique smallest ideal $J \subseteq R$ such that $f^{p^e-1} \in J^{[p^e]} = (j^{p^e} \mid j \in J)$ (see for example [BMS08, Definition 2.2]). In Section 5 we present some families of polynomials

2010 *Mathematics Subject Classification.* Primary 13A35; Secondary 13N10, 14B05.

Key words and phrases. Algorithm, Differential operator, Frobenius map, Prime characteristic.

*Partially supported by MTM2013-40775-P.

†Partially supported by NSF Grant DMS-1259142.

which have level one, and we give some examples. In Section 6 we focus on Elliptic Curves $\mathcal{C} \subseteq \mathbb{P}_{\mathbb{F}_p}^2$, where \mathbb{F}_p is the finite field with p elements. We prove the following characterization:

Theorem 1.1. Let $p \in \mathbb{Z}$ be a prime number and let $\mathcal{C} \subseteq \mathbb{P}_{\mathbb{F}_p}^2$ be an elliptic curve defined by a cubic $f(x, y, z) \in \mathbb{F}_p[x, y, z]$. Then

- (i) \mathcal{C} is ordinary if and only if f has level one.
- (ii) \mathcal{C} is supersingular if and only if f has level two.

All computations in this article are made using the computer software Macaulay2 [GS].

2. PRELIMINARIES

The goal of this section is to review the definitions, notations and results that we use throughout this paper. Unless otherwise specified, k will denote a perfect field of prime characteristic p . Under this assumption, it is known (see [Gro67, IV, Théorème 16.11.2]) that the ring of k -linear differential operators over $R = k[x_1, \dots, x_d]$ can be expressed in the following way:

$$\mathcal{D}_R := R \langle D_{x_i, t} \mid i = 1, \dots, d \text{ and } t \geq 1 \rangle, \quad \text{where } D_{x_i, t} := \frac{1}{t!} \frac{\partial^t}{\partial x_i^t}.$$

This allows us to regard \mathcal{D}_R as a filtered ring. Indeed, one has that

$$\mathcal{D}_R = \bigcup_{e \geq 0} \mathcal{D}_R^{(e)}, \quad \text{where } \mathcal{D}_R^{(e)} := R \langle D_{x_i, t} \mid i = 1, \dots, d \text{ and } 1 \leq t \leq p^e - 1 \rangle.$$

Moreover, it is shown by A. Yekutieli (see [Yek92, 1.4.9]) that $\mathcal{D}_R^{(e)} = \text{End}_{R^{p^e}}(R)$, hence the previous filtration does not depend on the choice of coordinates.

Now, we fix additional notation; given an $\alpha = (a_1, \dots, a_d) \in \mathbb{N}^d$ we shall use the following multi-index notation:

$$\mathbf{x}^\alpha := x_1^{a_1} \cdots x_d^{a_d}.$$

With this notation, we set $\|\mathbf{x}^\alpha\| := \max\{a_1, \dots, a_d\}$. By abuse of notation, we will sometimes also use $\|\alpha\|$ instead of $\|\mathbf{x}^\alpha\|$. For any polynomial $g \in k[x_1, \dots, x_d]$, we define

$$\|g\| := \max_{\mathbf{x}^\alpha \in \text{supp}(g)} \|\mathbf{x}^\alpha\|,$$

where if $g = \sum_{\alpha \in \mathbb{N}^d} g_\alpha \mathbf{x}^\alpha$ (such that $g_\alpha = 0$ for all but a finite number of terms) the support of g is defined as

$$\text{supp}(g) := \{\mathbf{x}^\alpha \in R \mid g_\alpha \neq 0\}.$$

Moreover, we also define $\deg(g)$ as the total degree of g . Finally, for any ideal $J \subseteq R$, $J^{[p^e]}$ will denote the ideal generated by all the p^e -th powers of elements in J , or equivalently the ideal generated by the p^e -th powers of any set of generators of J .

2.1. The ideal of p^e -th roots. Due to the central role which the ideal of p^e -th roots plays throughout this article, we review some well-known definitions and facts (cf. [AMBL05, page 465] and [BMS08, Definition 2.2]).

Definition 2.1. Given $g \in R$, we set $I_e(g)$ to be the smallest ideal $J \subseteq R$ such that $g \in J^{[p^e]}$.

Remark 2.2. Assume that k is perfect. In our assumptions, the ring R is a free R^{p^e} -module, with basis given by the monomials $\{\mathbf{x}^\alpha \mid \|\alpha\| \leq p^e - 1\}$. If we write

$$g = \sum_{0 \leq \|\alpha\| \leq p^e - 1} g_\alpha^{p^e} \mathbf{x}^\alpha,$$

then $I_e(g)$ is the ideal of R generated by all the elements g_α [BMS08, Proposition 2.5].

Remark 2.3. Notice that, if g is a homogeneous polynomial, then, for all $e \in \mathbb{N}$, $I_e(g)$ is a homogeneous ideal. Indeed, if we write $g = \sum_{0 \leq |\alpha| \leq p^e - 1} g_\alpha^{p^e} \mathbf{x}^\alpha$, then we can assume without loss of generality that every $g_\alpha^{p^e} \mathbf{x}^\alpha$ has degree equal to $\deg(g)$. But then g_α must be homogeneous of degree

$$\deg(g_\alpha) = \frac{\deg(g) - \deg(\mathbf{x}^\alpha)}{p^e}.$$

Since $I_e(g)$ is generated by the the elements g_α , it is a homogeneous ideal.

We have the following easy properties (see [AMBL05, Lemma 3.2 and Lemma 3.4] for details).

Proposition 2.4. Given $f \in R$ a non-zero polynomial, and given $e \geq 0$, the following statements hold.

- (i) $I_e(f) = I_{e+1}(f^p)$.
- (ii) $I_e(f^{p^e-1}) \supseteq I_{e+1}(f^{p^{e+1}-1})$.

Note that part (ii) of Proposition 2.4 produces the following decreasing chain of ideals:

$$(2.1.1) \quad R = I_0(f^{p^0-1}) \supseteq I_1(f^{p^1-1}) \supseteq I_2(f^{p^2-1}) \supseteq I_3(f^{p^3-1}) \supseteq \dots$$

It is shown in [AMBL05] that under our assumptions this chain stabilizes. The smallest integer $e \in \mathbb{N}$ where the chain stabilizes plays a central role in this paper. We summarize the facts that we will need in the following theorem. See [AMBL05, Proposition 3.5, and Theorem 3.7] for details and proofs.

Theorem 2.5. Let k be a perfect field of prime characteristic p , let $R = k[x_1, \dots, x_d]$, and let $f \in R \setminus \{0\}$. Define

$$e := \inf \left\{ s \geq 1 \mid I_{s-1}(f^{p^{s-1}-1}) = I_s(f^{p^s-1}) \right\}.$$

Then, the following assertions hold.

- (i) The chain of ideals (2.1.1) stabilizes rigidly, that is $e < \infty$ and $I_{e-1}(f^{p^{e-1}-1}) = I_{e+s}(f^{p^{e+s}-1})$ for any $s \geq 0$.
- (ii) One has

$$e = \min \left\{ s \geq 1 \mid f^{p^s-p} \in I_s(f^{p^s-1})^{[p^s]} \right\},$$

and $e \leq \deg(f)$.

- (iii) There exists $\delta \in \mathcal{D}_R^{(e)}$ such that $\delta(f^{p^e-1}) = f^{p^e-p}$, or equivalently such that $\delta(1/f) = 1/f^p$.
- (iv) There is no $\delta' \in \mathcal{D}_R^{(e')}$, with $e' < e$, such that $\delta'(1/f) = 1/f^p$.

Motivated by Theorem 2.5, we make the following definition.

Definition 2.6. For a non-zero polynomial $f \in R$, we call the integer e defined in Theorem 2.5 the *level of f* . Also, we will say that $\delta \in \mathcal{D}_R^{(e)}$ such that $\delta(f^{p^e-1}) = f^{p^e-p}$, or equivalently such that $\delta(1/f) = 1/f^p$, is a differential operator *associated with f* .

3. THE ALGORITHM

Let k be a computable perfect field of prime characteristic p (e.g., k is finite). Let $R = k[x_1, \dots, x_d]$, and let $f \in R$ be a non-zero polynomial. We now describe in details the algorithm that computes a differential operator $\delta \in \mathcal{D}_R$ associated with f .

- **Step 1.** Find the smallest integer $e \in \mathbb{N}$ such that $I_e(f^{p^e-p}) = I_e(f^{p^e-1})$. There is an implemented algorithm for the computation of the level of a given polynomial $f \in R$. Here follows a description:

Algorithm 3.1. Let k be a computable perfect field of prime characteristic p (e.g., finite), let $R = k[x_1, \dots, x_d]$, and let $f \in R$. These data act as the input of the procedure. Initialize $e = 0$ and $flag = true$. While $flag$ has the value $true$, execute the following commands:

- (i) Assign to e the value $e + 1$, and to q the value p^e .
 - (ii) Compute $I_e(f^{q-1})$.
 - (iii) Assign to J the value $I_e(f^{q-1})^{[q]}$.
 - (iv) If $f^{q-p} \in J$, then $flag = false$; otherwise, come back to step (i).
- At the end of this method, return the pair $(e, I_e(f^{p^e-1}))$. Such e is exactly the e described in Theorem 2.5, i.e., the level of f .

Remark 3.2. Since the level e is always at most $\deg(f)$, we can ensure that the While loop in Algorithm 3.1 finishes after, at most, $\deg(f)$ iterations. Notice that, a priori, there is a black box in this algorithm, namely the computation of $I_e(f^{q-1})$ at each step. However, the calculation of the ideal of p^e -th roots is well known (cf. [KS12a, Section 6]).

Remark 3.3. As pointed out by E. Canton in [Can15, Definition 2.3], the so-called *non-F-pure ideal* of f introduced by O. Fujino, K. Schwede and S. Takagi in [FST11, Definition 14.4] turns out to be $I_e(f^{p^e-1})$, where e is the level of f (see [FST11, Remark 16.2]). Therefore, Algorithm 3.1 provides a procedure to calculate the non-F-pure ideal.

- **Step 2.** For $e \in \mathbb{N}$ as in **Step 1** write $f^{p^e-1} = \sum_{i=1}^n c_i^{p^e} \mu_i$, where $\{\mu_1, \dots, \mu_n\}$ is the basis of R as an R^{p^e} -module consisting of all the monomials $x_1^{a_1} \cdots x_d^{a_d}$, with $a_i \leq p^e - 1$ for all $i = 1, \dots, d$.

Claim 3.4. For all $i = 1, \dots, n$ there exists $\delta_i \in \mathcal{D}_R^{(e)}$ such that $\delta_i(\mu_j) = 1$ if $i = j$ and $\delta_i(\mu_j) = 0$ if $i \neq j$.

Proof. For $i \in \{1, \dots, n\}$ and $\mu_i = x_1^{a_1} \cdots x_d^{a_d}$ consider $\nu_i := x_1^{p^e-1-a_1} \cdots x_d^{p^e-1-a_d}$, which is a monomial in R because $a_k \leq p^e - 1$ for all $k = 1, \dots, d$. Notice that $\nu_i \mu_j = (x_1 \cdots x_d)^{p^e-1}$ if and only if $i = j$. Then set

$$\delta_i := \left(\prod_{k=1}^d D_{x_k, p^e-1} \right) \cdot \nu_i \in \mathcal{D}_R^{(e)}.$$

Notice that $\delta_i(\mu_i) = 1$, and that if $\mu_j = x_1^{b_1} \cdots x_d^{b_d}$, then $\delta_i(\mu_j) = 0$ if $b_k < a_k$ for some $k \in \{1, \dots, d\}$. So let us assume that $a_k \leq b_k \leq p^e - 1$ for all $k = 1, \dots, d$, and that there is $s \in \{1, \dots, d\}$ such that $a_s < b_s$. Note that by definition of ν_i we have that $\nu_i \mu_j = x_1^{r_1} \cdots x_d^{r_d}$, with $p^e \leq r_s \leq 2p^e - 2$, so that we can write $r_s = p^e + n$ for some integer n with $0 \leq n \leq p^e - 2$.

Subclaim 3.5. The coefficient of $D_{x_s, p^e-1}(\nu_i \mu_j)$ is

$$\binom{p^e + n}{p^e - 1} \equiv 0 \pmod{p}.$$

Proof. As a consequence of a theorem proved by Lucas in [Luc78, pp. 51–52], it is enough to check that at least one of the digits of the base p expansion of $p^e - 1$ is greater than the corresponding digit in the base p expansion of $p^e + n$. The base p expansion of $p^e - 1$ is given by

$$p^e - 1 = (p - 1)(1 + p + \cdots + p^{e-1}) = (p - 1)p^0 + (p - 1)p^1 + \cdots + (p - 1)p^{e-1},$$

so that the subclaim is proved unless the first e digits of $p^e + n$ are $p - 1$ as well. But in this case, since $p^e + n > p^e - 1$ we would get

$$p^e + n \geq (p - 1)p^0 + (p - 1)p^1 + \cdots + (p - 1)p^{e-1} + p^e = 2p^e - 1,$$

a contradiction since $n \leq p^e - 2$. \square

The Subclaim shows that $D_{x_s, p^e-1}(\nu_i \mu_j) = 0$ for all μ_j with $j \neq i$. Therefore, using that $\delta_i \in \mathcal{D}_R^{(e)} = \text{End}_{R^{p^e}}(R)$, we get

$$\delta_i(f^{p^e-1}) = \delta_i \left(\sum_{j=1}^n c_j^{p^e} \mu_j \right) = \sum_{j=1}^n c_j^{p^e} \delta_i(\mu_j) = c_i^{p^e}. \quad \square$$

- **Step 3.** Since $1 \in \mathcal{D}_R^{(e)}$, for $e \in \mathbb{N}$ as in **Step 1** we have

$$f^{p^e-p} \in \mathcal{D}_R^{(e)}(f^{p^e-p}) = I_e(f^{p^e-p})^{[p^e]} = I_e(f^{p^e-1})^{[p^e]} = (c_1, \dots, c_n)^{[p^e]}.$$

In particular there exist $\alpha_1, \dots, \alpha_n \in R$ such that $f^{p^e-p} = \sum_{i=1}^n \alpha_i c_i^{p^e}$. Consider $\delta_i \in \mathcal{D}_R^{(e)}$ as in **Step 2**, so that $\delta_i(f^{p^e-1}) = c_i^{p^e}$, and set

$$\delta := \sum_{i=1}^n \alpha_i \delta_i \in \mathcal{D}_R^{(e)}.$$

With this choice we have

$$\delta(f^{p^e-1}) = \delta \left(\sum_{j=1}^n c_j^{p^e} \mu_j \right) = \sum_{i,j=1}^n c_j^{p^e} \alpha_i \delta_i(\mu_j) = \sum_{i=1}^n \alpha_i c_i^{p^e} = f^{p^e-p},$$

and using that $\delta \in \mathcal{D}_R^{(e)}$ we finally get

$$\delta \left(\frac{1}{f} \right) = \delta \left(\frac{f^{p^e-1}}{f^{p^e}} \right) = \frac{1}{f^{p^e}} \delta(f^{p^e-1}) = \frac{f^{p^e-p}}{f^{p^e}} = \frac{1}{f^p}.$$

4. THE MONOMIAL CASE

Throughout this section, let k be a perfect field and let $R = k[x_1, \dots, x_d]$. We now analyze the case when $f \in R$ is a monomial. First we show a lower bound for the level of f .

Lemma 4.1. Let $f = x_1^{a_1} \cdots x_d^{a_d}$ be a monomial in $R = k[x_1, \dots, x_d]$, with $a_i > 0$ for all $i = 1, \dots, d$. Let $\delta \in \mathcal{D}_R^{(e)}$ be such that $\delta(1/f) = 1/f^p$. Then, setting $a := \|f\| = \max\{a_i \mid 1 \leq i \leq d\}$, we have

$$e \geq \lceil \log_p(a) \rceil + 1.$$

Proof. It suffices to show that for $t := \lceil \log_p(a) \rceil$ we have $I_t(f^{p^t-p}) \not\supseteq I_t(f^{p^t-1})$. This is because if the chain

$$I_t(f^{p^t-p^{t-1}}) \supseteq I_t(f^{p^t-p^{t-2}}) \supseteq \dots \supseteq I_t(f^{p^t-p}) \supseteq I_t(f^{p^t-1})$$

stabilizes before such step, then it would be stable at it as well, and the smallest s such that $I_s(f^{p^s-p}) = I_s(f^{p^s-1})$ is precisely the level of f (see Theorem 2.5). Notice that $t = 0$ if and only if $a_i = 1$ for all i . For such a monomial the Lemma is trivially true. So let us assume that $t \geq 1$, or equivalently that $a_i \geq 2$ for at least one $i \in \{1, \dots, d\}$. Let

$$j_i := \min\{j \in \mathbb{N} \mid jp^t \geq a_i\},$$

and notice that $j_i \leq a_i$ for all i , and by choice of t we have that $j_i \geq 2$ for at least one i , say $j_1 \geq 2$. Then

$$f^{p^t-p} = x_1^{p^t a_1 - p a_1} \cdots x_d^{p^t a_d - p a_d} = (x_1^{a_1 - j_1} \cdots x_d^{a_d - j_d})^{p^t} \cdot x_1^{j_1 p^t - p a_1} \cdots x_d^{j_d p^t - p a_d}.$$

Since $(j_i - 1)p^t < a_i$ by definition of j_i , we have $j_i p^t - p a_i < p^t + a_i - p a_i < p^t$. This shows that $I_t(f^{p^t-p}) = (x_1^{a_1 - j_1} \cdots x_d^{a_d - j_d})$. On the other hand:

$$f^{p^t-1} = x_1^{p^t a_1 - a_1} \cdots x_d^{p^t a_d - a_d} = (x_1^{a_1 - 1} \cdots x_d^{a_d - 1})^{p^t} \cdot x_1^{p^t - a_1} \cdots x_d^{p^t - a_d},$$

which makes sense because $p^t \geq a_i$ for all i . This shows that $I_t(f^{p^t-1}) = (x_1^{a_1-1} \cdots x_d^{a_d-1})$, and because $j_1 \geq 2$ we have

$$I_t(f^{p^t-p}) = (x_1^{a_1-j_1} \cdots x_d^{a_d-j_d}) \supseteq (x_1^{a_1-2} \cdots x_d^{a_d-1}) \supsetneq (x_1^{a_1-1} \cdots x_d^{a_d-1}) = I_t(f^{p^t-1}). \quad \square$$

Theorem 4.2. Let $f = x_1^{a_1} \cdots x_d^{a_d}$ be a monomial in $k[x_1, \dots, x_d]$, with $a_i > 0$ for all $i = 1, \dots, d$. Let $a := \|f\| = \max\{a_i \mid 1 \leq i \leq d\}$. Then f has level $e := \lceil \log_p(a) \rceil + 1$, and $I_e(f^{p^e-1}) = (x_1^{a_1-1} \cdots x_d^{a_d-1})$. Furthermore,

$$\delta := \prod_{i=1}^d \left(x_i^{p^e - pa_i} \cdot D_{x_i, p^e-1} \cdot x_i^{a_i-1} \right) \in \mathcal{D}_R^{(e)}$$

is a differential operator associated with f .

Proof. Set $e := \lceil \log_p(a) \rceil + 1$. During the proof of Lemma 4.1 we already proved that $I_e(f^{p^e-1}) = (x_1^{a_1-1} \cdots x_d^{a_d-1})$; keeping this fact in mind, it is enough to check that $\delta(f^{p^e-1}) = f^{p^e-p}$. Indeed, we have

$$\begin{aligned} \delta(f^{p^e-1}) &= \delta \left(x_1^{p^e a_1 - a_1} \cdots x_d^{p^e a_d - a_d} \right) = \left(x_1^{a_1-1} \cdots x_d^{a_d-1} \right)^{p^e} \delta \left(x_1^{p^e - a_1} \cdots x_d^{p^e - a_d} \right) = \\ &= \prod_{i=1}^d \left(x_i^{p^e a_i - pa_i} \cdot D_{x_i, p^e-1} (x_i^{p^e-1}) \right) = x_1^{p^e a_1 - pa_1} \cdots x_d^{p^e a_d - pa_d} = f^{p^e-p}, \end{aligned}$$

and therefore the proof is completed. \square

Regarding the level e obtained in Theorem 4.2, one might ask whether, given any non-zero $f \in R$, its level would always be bounded above by $\lceil \log_p(\|f\|) \rceil + 1$. Unfortunately, this is not the case, as the following example illustrates.

Example 4.3. Consider $f := xy^3 + x^3 \in \mathbb{F}_2[x, y]$. In this case, one can check with Macaulay2 that the level of f is 4, while $\lceil \log_2(\|f\|) \rceil + 1 = 3$. In fact, the level is even strictly greater than $\lceil \log_2(\deg(f)) \rceil + 1 = 3$.

The monomial in Theorem 4.2 is assumed to be of the form $x_1^{a_1} \cdots x_d^{a_d}$. Using a suitable linear change of coordinates, we immediately get the following Corollary, which includes the general monomial case.

Corollary 4.4. Let $n \leq d$, let $f = \ell_1^{a_1} \cdots \ell_n^{a_n}$ be a product of powers of linear forms which are linearly independent over k , and let $a := \max\{a_i \mid 1 \leq i \leq n\}$. Then f has level $e = \lceil \log_p(a) \rceil + 1$, the ideal of p^e -th roots is $I_e(f^{p^e-1}) = (\ell_1^{a_1-1} \cdots \ell_n^{a_n-1})$ and

$$\delta := \prod_{i=1}^n \left(\ell_i^{p^e - pa_i} \cdot D_{\ell_i, p^e-1} \cdot \ell_i^{a_i-1} \right) \in \mathcal{D}_R^{(e)}$$

is a differential operator associated with f . Here, if $\ell_i = \sum_{j=1}^d \lambda_{ij} x_j$, then the differential operator D_{ℓ_i, p^e-1} is defined as $\sum_{j=1}^d \lambda_{ij} D_{x_j, p^e-1}$.

5. FAMILIES OF LEVEL ONE

Polynomials of level one, that is polynomials f such that $I_1(f^{p-1}) = R$, are somehow special. For instance, let $f, g \in R$ and let $\delta \in \mathcal{D}_R$ be associated with f . Assume that $e = 1$, then for $\delta' := \delta(g^{p-1} \cdot _)$ we get

$$\delta' \left(\frac{g}{f} \right) = \delta \left(\frac{g^p}{f} \right) = g^p \cdot \delta \left(\frac{1}{f} \right) = \left(\frac{g}{f} \right)^p.$$

The authors do not know whether, for any choice of $f, g \in R$, $f \neq 0$, there always exists $\delta' \in \mathcal{D}_R$ such that $\delta'(g/f) = (g/f)^p$. In fact, when $\delta \in \mathcal{D}_R^{(e)}$ for $e \geq 2$, the best we can get is $\delta'(g/f) = g^{p^e}/f^p$, with $\delta' := \delta(g^{p^{e-1}} \cdot _)$. On the other hand, for any $f \in R$ we have $R_f \cong \mathcal{D}_R \cdot \frac{1}{f}$ and therefore, for any $g \in R$, there exists $\delta' \in \mathcal{D}_R$ such that $\delta'(1/f) = g^p/f^p$. In fact it is enough to set $\delta' := g^p \cdot \delta$.

We will now exhibit some families of polynomials that have level one, together with some examples. However, before doing so, we want to single out the following elementary statement, because we will be using it repeatedly throughout this section. It may be regarded as a straightforward sufficient condition which ensures that a polynomial has level one. In this section, unless otherwise stated, k will denote a perfect field and $R = k[x_1, \dots, x_d]$ will be a polynomial ring over k .

Lemma 5.1. Let $f \in R$ be a non-zero polynomial, write

$$f^{p^e-1} = \sum_{0 \leq \|\alpha\| \leq p^e-1} f_\alpha^{p^e} \mathbf{x}^\alpha,$$

and assume that f_β is a unit for some $0 \leq \|\beta\| \leq p^e - 1$ and some $e \geq 1$. Then, f has level one.

Proof. By definition, we have that $I_e(f^{p^e-1}) = R$; on the other hand, we know that $I_1(f^{p-1}) \supseteq I_e(f^{p^e-1})$. In this way, combining these two facts it follows that $I_1(f^{p-1}) = R$, and therefore f has level one. \square

We can give an easy but useful characterization of homogeneous polynomials of level one.

Lemma 5.2. Let $f \in R$ be a homogeneous non-zero polynomial. Let $\{\mu_j\}_{j=1}^{p^d} := \{\mathbf{x}^\alpha \mid \|\alpha\| \leq p-1\}$ be the monomial basis of R as a R^p -module. Then f has level one if and only if $\mu_j \in \text{supp}(f^{p-1})$ for some $j = 1, \dots, p^d$.

Proof. Write

$$f^{p-1} = \sum_{0 \leq \|\alpha\| \leq p-1} f_\alpha^p \mathbf{x}^\alpha.$$

Note that, since f is homogeneous, $I_1(f^{p-1})$ is a homogeneous ideal by Remark 2.3. If f has level one, then $I_1(f^{p-1}) = (f_\alpha \mid 0 \leq \|\alpha\| \leq p-1) = R$, therefore there exists at least one coefficient f_β that is outside of the irrelevant maximal ideal $\mathfrak{m} = (x_1, \dots, x_d)$. Write $f_\beta = \lambda + r$, with $\lambda \in k$ and $r \in \mathfrak{m}$. Then, we can write $f^{p-1} = \lambda^p \mathbf{x}^\beta + h$ for some $h \in R$. Also, since $\{\mathbf{x}^\alpha \mid 0 \leq \|\alpha\| \leq p-1\}$ is a basis of R as R^p -module, there is no cancellation between $\lambda^p \mathbf{x}^\beta$ and h . Thus, $\mu_j = \mathbf{x}^\beta \in \text{supp}(f^{p-1})$. Conversely, if $\mu_j = \mathbf{x}^\beta \in \text{supp}(f^{p-1})$ for some $0 \leq \|\beta\| \leq p-1$, then we can write again $f^{p-1} = \lambda^p \mathbf{x}^\beta + h$ for some $\lambda \in k$ and some $h \in R$. Also, we can assume that $\mathbf{x}^\beta \notin \text{supp}(h)$. Then the coefficient f_β^p of \mathbf{x}^β in the expansion of f^{p-1} must be $\lambda^p + r^p = (\lambda + r)^p$ for some $r \in \mathfrak{m}$, and thus $\lambda + r \in I_1(f^{p-1})$. Since the latter is homogeneous (here we are using again Remark 2.3), we have in particular that $\lambda \in I_1(f^{p-1})$, which implies $I_1(f^{p-1}) = R$, and therefore f has level one. \square

Proposition 5.3. Let $f \in R$ be a non-zero polynomial whose support contains a squarefree term involving a variable that does not appear in any other term of the support of f . Then f has level one.

Proof. Without loss of generality we can assume that $x_1 \cdots x_n \in \text{supp}(f)$, and that x_1 does not appear in any other term of $\text{supp}(f)$. Write $f = \lambda_0 x_1 \cdots x_n + \sum_{i=1}^s \lambda_i m_i$, where $m_i = x^{\alpha_i}$ and $\alpha_i \in \mathbb{N}^d$ are of the form $(0, \alpha_{i2}, \dots, \alpha_{id})$ for $i = 1, \dots, s$. Then,

$$f^{p-1} = \sum_{i_0+i_1+\dots+i_s=p-1} \binom{p-1}{i_0, \dots, i_s} \lambda_0^{i_0} \lambda_1^{i_1} \cdots \lambda_s^{i_s} (x_1 \cdots x_n)^{i_0} m_1^{i_1} \cdots m_s^{i_s}.$$

Notice that in order for a term in the support of f^{p-1} to be divisible by x_1^{p-1} it is necessary that $i_0 = p-1$, in which case $i_1 = \dots = i_s = 0$. Hence $x_1^{p-1} \dots x_n^{p-1}$ is in the support of f^{p-1} , appearing with coefficient $\lambda_0^{p-1} \neq 0$. Then we can write

$$f^{p-1} = \lambda_0^{p-1} x_1^{p-1} \dots x_n^{p-1} + \sum_{\substack{0 \leq |\alpha| \leq p-1 \\ \alpha_1 \neq p-1}} f_\alpha^p \mathbf{x}^\alpha,$$

and the Proposition now follows from Lemma 5.1, since λ_0^{p-1} is a unit. \square

Example 5.4. Let $f = x^2 + y^2 + xyz \in \mathbb{F}_p[x, y, z]$. Since z appears in the square free term xyz and nowhere else in the support of f we have that f has level one by Proposition 5.3. In fact, $\delta := D_{x,p-1} D_{y,p-1} D_{z,p-1} \in \mathcal{D}_R^{(1)}$ is a differential operator associated with f .

Proposition 5.5. Let $f \in R$ be a non-zero polynomial of degree n such that every element of its support is a squarefree monomial. Then f has level one.

Proof. Without loss of generality we can assume that $x_1 \dots x_n \in \text{supp}(f)$. We want to show that we can apply Lemma 5.1. Let $f = \lambda_0 x_1 \dots x_n + \sum_{i=1}^s \lambda_i m_i$, where the monomials m_i are squarefree of degrees $d_i := \deg(m_i) \leq n$, that we can assume different from $x_1 \dots x_n$. Then

$$f^{p-1} = \sum_{i_0+i_1+\dots+i_s=p-1} \binom{p-1}{i_0, \dots, i_s} \lambda_0^{i_0} \lambda_1^{i_1} \dots \lambda_s^{i_s} (x_1 \dots x_n)^{i_0} m_1^{i_1} \dots m_s^{i_s}.$$

Note that the choice $i_0 = p-1, i_1 = \dots = i_s = 0$ gives the monomial $\lambda_0^{p-1} (x_1 \dots x_n)^{p-1}$, and we want to show that this choice of indices is the only one that gives such a monomial. By way of contradiction, assume that $(x_1 \dots x_n)^{i_0} m_1^{i_1} \dots m_s^{i_s} = (x_1 \dots x_n)^{p-1}$, then necessarily each m_i divides $x_1 \dots x_n$, because they are squarefree. Since we are assuming that none of the monomials m_i is equal to $x_1 \dots x_n$, we must have that $\deg(m_i) < n$. But then

$$\deg \left((x_1 \dots x_n)^{i_0} m_1^{i_1} \dots m_s^{i_s} \right) = n i_0 + d_1 i_1 + \dots + d_s i_s < n(i_0 + i_1 + \dots + i_s) = n(p-1),$$

which is a contradiction because $(x_1 \dots x_n)^{i_0} m_1^{i_1} \dots m_s^{i_s} = (x_1 \dots x_n)^{p-1}$, and the degree of the latter is $n(p-1)$. Therefore if we write

$$f^{p-1} = \sum_{0 \leq |\alpha| \leq p-1} f_\alpha^p \mathbf{x}^\alpha,$$

then the coefficient of $x_1^{p-1} \dots x_n^{p-1}$ is precisely λ_0^{p-1} , which is a unit. Using Lemma 5.1, the Proposition now follows. \square

Example 5.6. Let $R = \mathbb{F}_p[X_{ij}]_{1 \leq i, j \leq n}$ be a polynomial ring in n^2 variables and let $f = \det(X_{ij})$. Because of Proposition 5.5 f has level one, since its support consists only of squarefree monomials.

Proposition 5.7. Let $f \in R = k[x_1, \dots, x_d]$ be a homogeneous quadric. Then f has level one unless f is the square of a linear form, in which case f has level two.

Proof. If f is a power of a linear form, then f has level two by Corollary 4.4. Otherwise, if $p \neq 2$ there exists a linear change of variables that diagonalizes f (cf. [Ser73, Chapter IV, Proposition 5]). That is, we can assume that, after a linear change of coordinates, $f = x_1^2 + \dots + a x_n^2$, where $2 \leq n \leq d$ and a is either 1 or an element of k which is not a square. Notice that $x_1^{p-1} x_2^{p-1}$ appears with coefficient $\lambda := \binom{p-1}{\frac{p-1}{2}} \in k \setminus \{0\}$ if $n \geq 3$, and with coefficient $a^{(p-1)/2} \binom{p-1}{\frac{p-1}{2}} \in k \setminus \{0\}$ if $n = 2$. Therefore f has level one by Lemma 5.2. Finally, if $p = 2$ and f is not a power of a linear form, then we can assume that $x_1 x_2$ appears with non-zero coefficient in $f^{p-1} = f$, and we conclude using again Lemma 5.2. \square

Proposition 5.8. Let $f = x_1^t + \cdots + x_d^t \in R$ be a diagonal hypersurface of degree $t \geq 1$. If $t \leq \min\{d, p\}$ and $p \equiv 1 \pmod{t}$, then f has level one.

Proof. Our assumptions on t, d and p allow us to expand f^{p-1} in the following manner:

$$f^{p-1} = \frac{(p-1)!}{\left(\left(\frac{p-1}{t}\right)!\right)^t} x_1^{p-1} \cdots x_t^{p-1} + \sum_{\substack{i_1 + \cdots + i_d = p-1 \\ (i_1, \dots, i_t) \neq ((p-1)/t, \dots, (p-1)/t)}} \frac{(p-1)!}{i_1! \cdots i_d!} x_1^{i_1} \cdots x_d^{i_d}.$$

Since $\frac{(p-1)!}{\left(\left(\frac{p-1}{t}\right)!\right)^t} \in k \setminus \{0\}$, the above equality shows that $x_1^{p-1} \cdots x_t^{p-1}$ appears in $\text{supp}(f^{p-1})$ with non-zero coefficient, hence f has level one by Lemma 5.2. \square

The assumptions $p \equiv 1 \pmod{t}$ and $t \leq \min\{d, p\}$ in Proposition 5.8 cannot be removed in general, as the following examples illustrate.

Example 5.9. Let $R := \mathbb{F}_5[x, y, z]$ and $f := x^3 + y^3 + z^3$. One can check using Macaulay2 that f has level two. Notice that, in this case $3 = \deg(f) \leq \min\{3, 5\}$ and $5 \equiv 2 \pmod{3}$.

On the other hand, consider now $R := \mathbb{F}_7[x, y]$ and $f := x^3 + y^3$. One can check using Macaulay2 that f has level two. In this case $3 = \deg(f) > 2 = \min\{2, 7\}$ and $p = 7 \equiv 1 \pmod{3}$.

The diagonal hypersurface considered in Proposition 5.8 is of the form $x_1^t + \cdots + x_d^t$; using a suitable linear change of coordinates, we immediately get the following Corollary, which includes as a particular case Proposition 5.8.

Corollary 5.10. Let $n \leq d$, let $f = \ell_1^t + \cdots + \ell_n^t$ be a diagonal hypersurface of degree $t \geq 1$ made up by linear forms ℓ_1, \dots, ℓ_n which are linearly independent over the field k . If $t \leq \min\{n, p\}$ and $p \equiv 1 \pmod{t}$, then f has level one.

Before going on, we want to review the following notion (see [ILL⁺07, page 243]):

Definition 5.11. A polynomial $f \in R$ is said to be *regular* provided

$$\text{Tj}(f) := \left(f, \frac{\partial f}{\partial x_1}, \dots, \frac{\partial f}{\partial x_d}\right) = R,$$

where $\text{Tj}(f)$ denotes the Tjurina ideal attached to f .

In characteristic zero, a polynomial is regular if and only if its Bernstein-Sato polynomial is $b_f(s) = s + 1$ [ILL⁺07, Theorem 23.12]. In this case, R_f is generated by $1/f$ as a D -module.

Proposition 5.12. Let k be a perfect field of characteristic 2, and let $f \in k[x_1, \dots, x_d]$ be regular. Then, f has level one.

Proof. Since f is regular, there are $r_0, r_1, \dots, r_d \in R$ such that

$$1 = r_0 f + r_1 \frac{\partial f}{\partial x_1} + \cdots + r_d \frac{\partial f}{\partial x_d}.$$

In this way, setting

$$\delta := r_0 + \sum_{j=1}^d \frac{\partial}{\partial x_j}$$

it follows that $\delta(f) = 1$ and therefore f has level one. \square

Remark 5.13. A very easy way to produce polynomials which are simultaneously regular and of level one in arbitrary prime characteristic works as follows. Let k be a perfect field of prime characteristic p , let $R = k[x_1, \dots, x_d]$, and assume that $f \in R$ is a non-zero polynomial of the form $f = \lambda x_i + g$, for some $1 \leq i \leq d$, some $\lambda \in k - \{0\}$, and some $g \in R$ such that

$$\frac{\partial g}{\partial x_i} = 0 \quad (\text{i.e., } g \in k[x_1, \dots, x_{i-1}, \widehat{x_i}, x_{i+1}, x_{i+2}, \dots, x_d]).$$

Then, f is regular and of level one; indeed, the fact that f is of level one follows directly from Proposition 5.3.

6. ELLIPTIC CURVES

Let $p \in \mathbb{Z}$ be a prime and let $\mathcal{C} \subseteq \mathbb{P}_{\mathbb{F}_p}^2$ be an elliptic curve defined by an homogeneous cubic $f(x, y, z) \in \mathbb{F}_p[x, y, z]$. We want to review here the following notion (see [Hus87, 13.3.1]).

Definition 6.1. \mathcal{C} is said to be ordinary if the monomial $(xyz)^{p-1}$ appears in the expansion of f^{p-1} with non-zero coefficient. Otherwise, \mathcal{C} is said to be supersingular.

The general form of a cubic defining an elliptic curve is the following

$$f = y^2z + a_1xyz + a_3yz^2 - x^3 - a_2x^2z - a_4xz^2 - a_6z^3,$$

where $a_1, \dots, a_6 \in \mathbb{F}_p$. When $p \neq 2, 3$, the expression above can be further simplified to

$$f = y^2z - x^3 + axz^2 + bz^3,$$

for $a, b \in \mathbb{F}_p$ (see [Hus87, 3.3.6] for details). We are now interested in computing the level of elliptic curves \mathcal{C} . We are mainly interested in upper bounds, since it is easy to see from Lemma 5.2 that any ordinary elliptic curve has level one, and that any supersingular elliptic curve has level at least two. First, we explore the low characteristic cases, where the list of possibilities (up to isomorphism) is very limited.

Proposition 6.2. Let $\mathcal{C} \subseteq \mathbb{P}_{\mathbb{F}_p}^2$ be a supersingular elliptic curve defined by a cubic $f \in \mathbb{F}_p[x, y, z]$. If $p = 2$ or $p = 3$, then f has level two.

Proof. Set $D := D_{x,p^2-1}D_{y,p^2-1}D_{z,p^2-1}$. By [Hus87, 13.3.2 and 13.3.3] there are only the following two cases, up to isomorphism:

| p | Elliptic curve | Differential operator |
|-----|---------------------|---|
| 2 | $x^3 + y^2z + yz^2$ | $y^2Dx^3z + z^2Dx^3y + x^2Dxyz^2$ |
| 3 | $x^3 - xz^2 - y^2z$ | $(x^6z^3 - x^3y^6)Dx^4z^5 +$ $+(x^9 + x^3z^6 + y^6z^3)Dxy^8 + y^3z^6Dx^4y^5$ |

The table above is exhibiting a differential operator of level two for each polynomial, therefore the level is at most two in all such cases. We have already noticed that \mathcal{C} is ordinary if and only if f has level one. This shows that when $p = 2$ or $p = 3$, \mathcal{C} is supersingular if and only if f has level two. \square

Recall that $R = \mathbb{F}_p[x, y, z]$ is a free R^{p^2} -module with basis given by $\{x^r y^s z^t \mid 0 \leq r, s, t \leq p^2 - 1\}$. For a polynomial $g \in \mathbb{F}_p[x, y, z]$ consider

$$g = \sum_{0 \leq r, s, t \leq p^2 - 1} (c(r, s, t))^{p^2} x^r y^s z^t,$$

and recall that by Proposition 2.4, with this notation, $I_2(g)$ is the ideal generated by the elements $c(r, s, t)$, for $0 \leq r, s, t \leq p^2 - 1$.

Remark 6.3. If $f \in \mathbb{F}_p[x, y, z]$ is a cubic and $g = f^{p^2-1}$, then for any $0 \leq r, s, t \leq p^2 - 1$ one has

$$\deg(c(r, s, t)) \leq \left\lfloor \frac{\deg(f^{p^2-1})}{p^2} \right\rfloor = \left\lfloor \frac{3(p^2 - 1)}{p^2} \right\rfloor = 2.$$

In particular, $I_2(f^{p^2-1}) = (c(r, s, t) \mid 0 \leq r, s, t \leq p^2 - 1)$ is generated in degree at most two.

For the rest of the section, we will denote $I_1(f^{p^2-1})$ and $I_2(f^{p^2-1})$ simply by I_1 and I_2 .

6.1. Preliminary computations. The purpose of this part is to single out some technical facts which will be used for proving the main result of this section; namely, Theorem 6.9.

Lemma 6.4. Let $f = y^2z - x^3 + axz^2 + bz^3 \in \mathbb{F}_p[x, y, z]$, where $p \neq 2, 3$. Then

$$c(0, p^2 - 2, p^2 - 1) = y.$$

Proof. A monomial in the expansion of f^{p^2-1} will have the form $(y^2z)^h(-x^3)^i(axz^2)^j(bz^3)^k$, where $h + i + j + k = p^2 - 1$. Looking at the coefficient of $y^{p^2-2}z^{p^2-1}$ in such expansion, by degree considerations we only have three possibilities:

$$y^{2h}x^{3i+j}z^{h+2j+3k} = \begin{cases} x^{p^2} \cdot y^{p^2-2}z^{p^2-1} \\ y^{p^2} \cdot y^{p^2-2}z^{p^2-1} \\ z^{p^2} \cdot y^{p^2-2}z^{p^2-1} \end{cases}$$

Since $p^2 - 2$ is not even, there is no choice of h that realizes the first and the third cases. So we are left with the second, which is achieved only by the choice $h = p^2 - 1, i = j = k = 0$. This shows that the coefficient of $y^{p^2-2}z^{p^2-1}$ in the expansion of f^{p^2-1} is precisely y^{p^2} , and the Lemma follows. \square

Before going on, we need to review the following classical result, due to Legendre, because it will play some role later in this section (see Proof of Lemma 6.7). We refer to [AZ04, page 8] for a proof.

Theorem 6.5 (Legendre). Let $n \geq 0$ be a non-negative integer, let p be a prime number, and let $\sigma_p(n)$ be the sum of the base p digits of n . Then,

$$v_p(n!) = \frac{n - \sigma_p(n)}{p - 1},$$

where, given any non-negative integer $m \geq 0$,

$$v_p(m) := \max\{t \geq 0 : p^t \mid m\}.$$

Lemma 6.6. Let $p \neq 2$ be a prime. Then,

$$\lambda := \frac{(p^2 - 1)!}{\left(\left(\frac{p^2 - 1}{2}\right)!\right)^2} \not\equiv 0 \pmod{p}.$$

Proof. On one hand, $p^2 - 1 = (p - 1)(1 + p)$ is the base p expansion of $p^2 - 1$; on the other hand, since $p \neq 2$ it follows that

$$\frac{p^2 - 1}{2} = \left(\frac{p - 1}{2}\right)(1 + p)$$

is the base p expansion of $(p^2 - 1)/2$. Keeping in mind these two facts it follows, using Legendre's Theorem, that

$$\begin{aligned} v_p(\lambda) &= v_p((p^2 - 1)!) - 2v_p\left(\left(\frac{p^2 - 1}{2}\right)!\right) \\ &= \frac{p^2 - 1 - 2(p - 1)}{p - 1} - 2 \cdot \left(\frac{\frac{p^2 - 1}{2} - 2\left(\frac{p - 1}{2}\right)}{p - 1}\right) = p - 1 - (p - 1) = 0, \end{aligned}$$

hence p does not divide λ and therefore we can ensure that $\lambda \not\equiv 0 \pmod{p}$. \square

Lemma 6.7. Let $p \neq 2, 3$ be a prime, and let $f = y^2z - x^3 \in \mathbb{F}_p[x, y, z]$. Then, $I_1 = I_2 = (x, y)$. In particular, f has level two.

Proof. Consider the expansion

$$f^{p^2-1} = \sum_{i=0}^{p^2-1} \frac{(p^2-1)!}{i!(p^2-1-i)!} y^{2i} z^i x^{3(p^2-1-i)}.$$

For $i = (p^2-1)/2$ we obtain the monomial

$$\lambda y^{p^2-1} z^{(p^2-1)/2} x^{3(p^2-1)/2} = \lambda x^{p^2} \cdot \left(x^{(p^2-3)/2} y^{p^2-1} z^{(p^2-1)/2} \right),$$

where $\lambda = \frac{(p^2-1)!}{\left(\left(\frac{p^2-1}{2}\right)!\right)^2} \neq 0$ (indeed, this follows by Lemma 6.6). Because of the term z^i in the expansion above, the choice $i = (p^2-1)/2$ is clearly the only one that gives the monomial $x^{(p^2-3)/2} y^{p^2-1} z^{(p^2-1)/2}$ of the basis of R as an R^{p^2} -module. Therefore $c\left(\frac{p^2-3}{2}, p^2-1, \frac{p^2-1}{2}\right) = \lambda^{1/p^2} x = \lambda x$, and thus $x \in I_2$. In addition, by Lemma 6.4 we always have $y \in I_2$. Therefore $(x, y) \subseteq I_2$. On the other hand, consider the expansion

$$f^{p-1} = \sum_{j=0}^{p-1} \frac{(p-1)!}{j!(p-1-j)!} y^{2j} z^j x^{3(p-1-j)}.$$

We claim that either $2j \geq p$ or $3(p-1-j) \geq p$. In fact, suppose $j < p/2$, or equivalently $j \leq (p-1)/2$, because j is an integer. Then $3(p-1-j) \geq 3(p-1)/2 \geq p$ since $p \geq 5$ by assumption. This shows that all the coefficients $c(r, s, t)^p$ in the expansion of $f^{p-1} = \sum_{0 \leq r, s, t \leq p-1} c(r, s, t)^p x^r y^s z^t$ are contained in $(x, y)^{[p]}$, and thus $I_1 = (c(r, s, t) \mid 0 \leq r, s, t \leq p-1) \subseteq (x, y)$. Therefore the Lemma follows from the chain of inclusions $(x, y) \subseteq I_2 \subseteq I_1 \subseteq (x, y)$. \square

Lemma 6.8. Let $p \neq 2, 3$ be a prime and let \mathcal{C} be a supersingular elliptic curve defined by $f(x, y, z) = y^2z - x^3 + axz^2 + bz^3$. If either $a = 0$ or $b = 0$, then f has level two.

Proof. Notice that, since \mathcal{C} is supersingular, we have $\mathbb{F}_p[x, y, z] = R \supsetneq I_1 \supseteq I_2$, and we want to show that $I_1 = I_2$. Also, by Lemma 6.4 we always have $y \in I_2$. If $a = b = 0$, then Lemma 6.7 ensures that $I_1 = I_2 = (x, y)$. Now assume $a \neq 0$ and $b = 0$. We claim that $c\left(\frac{p^2-1}{2}, p^2-1, \frac{p^2-3}{2}\right) = \mu z$, for some $\mu \neq 0$. A monomial in the expansion of f^{p^2-1} will have the form $(y^2z)^h (-x^3)^i (axz^2)^j$ for $h + i + j = p^2-1$. Looking at the terms involving $x^{\frac{p^2-1}{2}} y^{p^2-1} z^{\frac{p^2-3}{2}}$ the possibilities are

$$y^{2h} x^{3i+j} z^{2j+h} = \begin{cases} x^{p^2} \cdot x^{\frac{p^2-1}{2}} y^{p^2-1} z^{\frac{p^2-3}{2}} \\ y^{p^2} \cdot x^{\frac{p^2-1}{2}} y^{p^2-1} z^{\frac{p^2-3}{2}} \\ z^{p^2} \cdot x^{\frac{p^2-1}{2}} y^{p^2-1} z^{\frac{p^2-3}{2}} \end{cases}$$

The second case is not possible, since $2p^2-1$ is not even and hence here is no h that gives it. This forces $h = (p^2-1)/2$, and hence the first case is also not possible, since for such an h the exponent of z must be at least $(p^2-1)/2$. Canceling $y^{p^2-1} z^{\frac{p^2-1}{2}}$ we are left with

$$x^{3i} z^{2j} = x^{\frac{p^2-1}{2}} z^{p^2-1}$$

This implies that $j = \frac{p^2-1}{2}$ and $i = 0$. Therefore the coefficient of $x^{\frac{p^2-1}{2}} y^{p^2-1} z^{\frac{p^2-3}{2}}$ in the expansion of f^{p^2-1} is

$$c\left(\frac{p^2-1}{2}, p^2-1, \frac{p^2-3}{2}\right)^{p^2} = a^{\frac{p^2-1}{2}} z^{p^2} = (\mu z)^{p^2},$$

where $\mu = \left(a^{\frac{p^2-1}{2}}\right)^{1/p^2} \neq 0$. This shows the claim. With similar considerations, one can see that

$$c\left(\frac{p^2-3}{2}, p^2-1, \frac{p^2-1}{2}\right) = x.$$

This gives $(x, y, z) \subseteq I_2 \subseteq I_1 \subsetneq R$, which forces $I_1 = I_2$ because I_1 is a proper homogeneous ideal of $R = \mathbb{F}_p[x, y, z]$, hence $I_1 \subseteq (x, y, z)$. Finally, if $a = 0$ and $b \neq 0$ the arguments are completely analogous to the case $a \neq 0, b = 0$. Here we get

$$c\left(\frac{p^2-3}{2}, p^2-1, \frac{p^2-1}{2}\right) = x \quad \text{and} \quad c(0, p^2-1, p^2-2) = \lambda z$$

for some $\lambda \neq 0$. Therefore $(x, y, z) \subseteq I_2 \subseteq I_1 \subsetneq R$, which once again forces $I_1 = I_2$. \square

6.2. Main result. Next statement is the main result of this section.

Theorem 6.9. Let $p \in \mathbb{Z}$ be a prime number and let $\mathcal{C} \subseteq \mathbb{P}_{\mathbb{F}_p}^2$ be an elliptic curve defined by a cubic $f(x, y, z) \in \mathbb{F}_p[x, y, z]$. Then

- (i) \mathcal{C} is ordinary if and only if f has level one.
- (ii) \mathcal{C} is supersingular if and only if f has level two.

Proof. Part (i) follows from Lemma 5.2, which also shows that if f has level at least two, then \mathcal{C} is supersingular. So it is left to show that if \mathcal{C} is supersingular, then f has level precisely equal to two. By Proposition 6.2 we only have to consider the cases where $p \neq 2, 3$, thus we can assume that f is of the form

$$f(x, y, z) = y^2z - x^3 + axz^2 + bz^3.$$

for $a, b \in \mathbb{F}_p$. Furthermore, by Lemma 6.8 we can assume that $ab \neq 0$. First we claim that $c\left(\frac{p^2-3}{2}, p^2-1, \frac{p^2-1}{2}\right) = x + \lambda z$, where

$$\lambda = \sum_{i=0}^{\frac{p^2-7}{6}} \frac{(p^2-1)!}{\left(\frac{p^2-1}{2}\right)! i! \left(\frac{p^2-3}{2} - 3i\right)! (2i+1)!} (-1)^i a^{\frac{p^2-3}{2} - 3i} b^{2i+1} \in \mathbb{F}_p.$$

In fact a general monomial in the expansion of f^{p^2-1} will have the form $(y^2z)^h (-x^3)^i (axz^2)^j (bz^3)^k$, and by looking at terms that involve $x^{\frac{p^2-3}{2}} y^{p^2-1} z^{\frac{p^2-1}{2}}$, by degree considerations we have three possibilities:

$$y^{2h} x^{3i+j} z^{h+2j+3k} = \begin{cases} x^{p^2} \cdot x^{\frac{p^2-3}{2}} y^{p^2-1} z^{\frac{p^2-1}{2}} \\ y^{p^2} \cdot x^{\frac{p^2-3}{2}} y^{p^2-1} z^{\frac{p^2-1}{2}} \\ z^{p^2} \cdot x^{\frac{p^2-3}{2}} y^{p^2-1} z^{\frac{p^2-1}{2}} \end{cases}$$

The second case cannot be realized since $2p^2 - 1$ is not even, hence we necessarily have $h = \frac{p^2-1}{2}$. This leaves two cases:

$$x^{3i+j} z^{2j+3k} = \begin{cases} x^{p^2} \cdot x^{\frac{p^2-3}{2}} \\ z^{p^2} \cdot x^{\frac{p^2-3}{2}} \end{cases}$$

The first one can only happen when $h = i = \frac{p^2-1}{2}$ and $j = k = 0$, giving the monomial

$$(y^2z)^{\frac{p^2-1}{2}} (-x^3)^{\frac{p^2-1}{2}} = (-1)^{\frac{p^2-1}{2}} x^{p^2} \cdot x^{\frac{p^2-3}{2}} y^{p^2-1} z^{\frac{p^2-1}{2}} = x^{p^2} \cdot x^{\frac{p^2-3}{2}} y^{p^2-1} z^{\frac{p^2-1}{2}},$$

because $p^2 \equiv 1 \pmod{4}$. For the second case i, j and k will have to satisfy

$$\begin{cases} j = \frac{p^2-3}{2} - 3i \\ k = 2i + 1 \end{cases}$$

As both j and k must be non-negative, we have

$$0 \leq i \leq \left\lfloor \frac{p^2-3}{6} \right\rfloor = \frac{p^2-7}{6},$$

because $p^2 \equiv 1 \pmod{6}$. Furthermore, the coefficient of $(y^2z)^h(-x^3)^i(axz^2)^j(bz^3)^k$ under these conditions is precisely

$$\frac{(p^2-1)!}{\left(\frac{p^2-1}{2}\right)!i!\left(\frac{p^2-3}{2}-3i\right)!(2i+1)!}(-1)^i a^{\frac{p^2-3}{2}-3i} b^{2i+1},$$

proving that

$$c\left(\frac{p^2-3}{2}, p^2-1, \frac{p^2-1}{2}\right)^{p^2} = x^{p^2} + \lambda z^{p^2} = (x + \lambda^{1/p^2} z)^{p^2},$$

for λ as above. Since we are working over \mathbb{F}_p , we finally have $\lambda^{1/p^2} = \lambda$, showing the claim. With similar arguments, one can see that

$$c\left(\frac{p^2-1}{2}, p^2-3, \frac{p^2+1}{2}\right) = ax + \mu z,$$

$$c\left(\frac{p^2-7}{2}, p^2-1, \frac{p^2+3}{2}\right) = -ax + \tau z,$$

$$c(0, p^2-1, p^2-2) = \theta x + b^{\frac{p^2-1}{2}} z,$$

where $\theta \in \mathbb{F}_p$,

$$\mu = \sum_{i=0}^{\frac{p^2-1}{6}} \frac{(p^2-1)!}{\left(\frac{p^2-3}{2}\right)!i!\left(\frac{p^2-1}{2}-3i\right)!(2i+1)!}(-1)^i a^{\frac{p^2-1}{2}-3i} b^{2i+1} \in \mathbb{F}_p,$$

and

$$\tau = \sum_{i=0}^{\frac{p^2-7}{6}} \frac{(p^2-1)!}{\left(\frac{p^2-1}{2}\right)!i!\left(\frac{p^2-7}{2}-3i\right)!(2i+3)!}(-1)^i a^{\frac{p^2-7}{2}-3i} b^{2i+3} \in \mathbb{F}_p.$$

Since $I_2 = (c(r, s, t) \mid 0 \leq r, s, t \leq p^2-1)$, in particular we have

$$(x + \lambda z, ax + \mu z, -ax + \tau z, \theta x + b^{\frac{p^2-1}{2}} z) \subseteq I_2.$$

Claim. The following matrix has full rank:

$$\begin{bmatrix} a & \mu \\ 1 & \lambda \\ -a & \tau \\ \theta & b^{\frac{p^2-1}{2}} \end{bmatrix}.$$

In fact if $\det \begin{bmatrix} a & \mu \\ 1 & \lambda \end{bmatrix} \neq 0$ then we are done, otherwise we have $\mu = a\lambda$. Note that

$$\begin{aligned} \mu &= \sum_{i=0}^{\frac{p^2-1}{6}} \frac{(p^2-1)!}{\left(\frac{p^2-3}{2}\right)!i! \left(\frac{p^2-1}{2}-3i\right)!(2i+1)!} (-1)^i a^{\frac{p^2-1}{2}-3i} b^{2i+1} \\ &= \left[\sum_{i=0}^{\frac{p^2-7}{6}} \frac{(p^2-1)!}{\left(\frac{p^2-3}{2}\right)!i! \left(\frac{p^2-1}{2}-3i\right)!(2i+1)!} (-1)^i a^{\frac{p^2-1}{2}-3i} b^{2i+1} \right] + (-1)^{\frac{p^2-1}{6}} \frac{(p^2-1)!}{\left(\frac{p^2-3}{2}\right)! \left(\frac{p^2-1}{6}\right)! \left(\frac{p^2+2}{3}\right)!} b^{\frac{p^2+2}{3}} \\ &= \left[\sum_{i=0}^{\frac{p^2-7}{6}} \frac{(p^2-1)!}{\left(\frac{p^2-3}{2}\right)!i! \left(\frac{p^2-1}{2}-3i\right)!(2i+1)!} (-1)^i a^{\frac{p^2-1}{2}-3i} b^{2i+1} \right] + \frac{3(p^2-1)!}{\left(\frac{p^2-1}{2}\right)! \left(\frac{p^2-7}{6}\right)! \left(\frac{p^2+2}{3}\right)!} b^{\frac{p^2+2}{3}}, \end{aligned}$$

where the last equality comes from the fact that $p^2 \equiv 1 \pmod{12}$ and by rearranging the binomial coefficients. Also,

$$a\lambda = \sum_{i=0}^{\frac{p^2-7}{6}} \frac{(p^2-1)!}{\left(\frac{p^2-1}{2}\right)!i! \left(\frac{p^2-3}{2}-3i\right)!(2i+1)!} (-1)^i a^{\frac{p^2-1}{2}-3i} b^{2i+1}.$$

Using that $\mu = a\lambda$, we get

$$\begin{aligned} &\frac{3(p^2-1)!}{\left(\frac{p^2-1}{2}\right)! \left(\frac{p^2-7}{6}\right)! \left(\frac{p^2+2}{3}\right)!} b^{\frac{p^2+2}{3}} \\ &= \sum_{i=1}^{\frac{p^2-7}{6}} \left[\frac{1}{\frac{p^2-1}{2}} - \frac{1}{\frac{p^2-1}{2}-3i} \right] \frac{(p^2-1)!}{\left(\frac{p^2-3}{2}\right)!i! \left(\frac{p^2-3}{2}-3i\right)!(2i+1)!} (-1)^i a^{\frac{p^2-1}{2}-3i} b^{2i+1} \\ &= \sum_{i=1}^{\frac{p^2-7}{6}} \frac{-3(p^2-1)!}{\left(\frac{p^2-1}{2}\right)!(i-1)! \left(\frac{p^2-1}{2}-3i\right)!(2i+1)!} (-1)^i a^{\frac{p^2-1}{2}-3i} b^{2i+1} \\ &= 3 \sum_{i=0}^{\frac{p^2-13}{6}} \frac{(p^2-1)!}{\left(\frac{p^2-1}{2}\right)!i! \left(\frac{p^2-7}{2}-3i\right)!(2i+3)!} (-1)^i a^{\frac{p^2-7}{2}-3i} b^{2i+3}, \end{aligned}$$

where the last equality comes from reindexing the sum. Since 3 is invertible in \mathbb{F}_p we get that

$$\begin{aligned} 0 &= \left[\sum_{i=0}^{\frac{p^2-13}{6}} \frac{(p^2-1)!}{\left(\frac{p^2-1}{2}\right)!i! \left(\frac{p^2-7}{2}-3i\right)!(2i+3)!} (-1)^i a^{\frac{p^2-7}{2}-3i} b^{2i+3} \right] - \left[\frac{(p^2-1)!}{\left(\frac{p^2-1}{2}\right)! \left(\frac{p^2-7}{6}\right)! \left(\frac{p^2+2}{3}\right)!} b^{\frac{p^2+2}{3}} \right] \\ &= \sum_{i=0}^{\frac{p^2-7}{6}} \frac{(p^2-1)!}{\left(\frac{p^2-1}{2}\right)!i! \left(\frac{p^2-7}{2}-3i\right)!(2i+3)!} (-1)^i a^{\frac{p^2-7}{2}-3i} b^{2i+3} \end{aligned}$$

because $p^2 \equiv 1 \pmod{12}$, hence $(-1)^{\frac{p^2-7}{6}} = -1$. But the latter is precisely τ , and this argument shows that if $a\lambda = \mu$, then $\tau = 0$. Therefore either $a\lambda \neq \mu$ or

$$\det \begin{bmatrix} -a & \tau \\ \theta & b^{\frac{p^2-1}{2}} \end{bmatrix} = \det \begin{bmatrix} -a & 0 \\ \theta & b^{\frac{p^2-1}{2}} \end{bmatrix} = -ab^{\frac{p^2-1}{2}} \neq 0.$$

Hence the matrix has rank two, and the Claim follows.

But this shows that there are linear combinations of $x + \lambda z$, $ax + \mu z$, $-ax + \tau z$ and $\theta x + b^{\frac{p^2-1}{2}} z$ that produce x and z , that is $(x, z) \subseteq I_2$. By Lemma 6.4 we always have that $y \in I_2$. Therefore

$$(x, y, z) \subseteq I_2 \subseteq I_1 \subsetneq R,$$

implying that $I_1 = I_2$ and hence that f has level two. \square

7. A MACAULAY2 SESSION

The purpose of this section is to explain, through a Macaulay2 session, how the algorithm introduced in Section 3 works in specific examples.

We begin clearing the previous input and loading our scripts.

```
clearAll;
```

```
load "differentialOperator.m2";
```

We fix the polynomial ring that we will use throughout the following examples.

```
p=2;
```

```
F=ZZ/p;
```

```
R=F[x,y,z,w];
```

The first example illustrates a particular case of Theorem 4.2.

```
i6 : f=x^3*y^5*z^7*w^4;
```

```
i7 : L=differentialOperatorLevel(f);
```

```
i8 : L
```

```
o8 = (4, ideal(x y z w ))
```

This means that, in this case, f has level 4 and that $I_4(f^{2^4-1}) = (x^2y^4z^6w^3)$. Now, we produce a differential operator δ of level 4 such that $\delta(1/f) = 1/f^2$.

```
i7 : DifferentialOperator(f)
```

```
o7 = | x10y6z2w8 d_0^15d_1^15d_2^15d_3^15x2y4z6w3 |
```

As the reader will note, the output is a row matrix; it means that δ turns out to be

$$x^{10}y^6z^2w^8 \cdot D_{x,2^4-1}D_{y,2^4-1}D_{z,2^4-1}D_{w,2^4-1} \cdot x^2y^4z^6w^3.$$

Our next aim is to illustrate a particular case of Corollary 4.4.

```
ii8 : f=x^3*(x+y)^5*(x+y+z)^7*(x+y+z+w)^4;
```

```
ii9 : L=differentialOperatorLevel(f);
```

```
ii10 : first L
```

```
oo10 = 4
```

Now, a particular case of Proposition 5.3.

```
ii13 : f=x^2+y^2+z^3+x*y*z*w;
```

```
ii14 : L=differentialOperatorLevel(f);
```

```
ii15 : L
```

```
oo15 = (1, ideal 1)
```

This means that f has level one. Now, we produce the corresponding differential operator.

```
i7 : DifferentialOperator(f)
```

```
o7 = | 1 d_0d_1d_2d_3 |
```

It means that the differential operator produced in this case is $D_{x,1}D_{y,1}D_{z,1}D_{w,1}$.

The following computation may be regarded as a particular case of Proposition 5.5.

```
i6 : f=x*w-y*z;
```

```
i7 : DifferentialOperator(f)
```

```
o7 = | 1 d_0d_1d_2d_3yz |
```

It means that, in this case, the differential operator produced is $D_{x,1}D_{y,1}D_{z,1}D_{w,1} \cdot yz$.

Next, a homogeneous quadric (cf. Proposition 5.7).

```
i6 : f=x^2+y^2+x*y+z^2+w^2;
```

```
i7 : DifferentialOperator(f)
```

```
o7 = | 1 d_0d_1d_2d_3zw |
```

We finish with a homogeneous cubic.

```
ii12 : f=x^3+y^3+z^3+w^3;
```

```
ii13 : L=differentialOperatorLevel(f);
```

```
ii14 : L
```

```
oo14 = (2, ideal (w, z, y, x))
```

```
oo14 : Sequence
```

```
ii15 : DifferentialOperator(f)
```

```
oo15 = | w2 d_0^3d_1^3d_2^3d_3^3x3z3w |
      | z2 d_0^3d_1^3d_2^3d_3^3x3zw3 |
      | y2 d_0^3d_1^3d_2^3d_3^3yz3w3 |
      | x2 d_0^3d_1^3d_2^3d_3^3xy3z3 |
```

This means that our differential operator in this case turns out to be

$$\begin{aligned} & w^2 \cdot D_{x,3}D_{y,3}D_{z,3}D_{w,3} \cdot x^3z^3w + z^2 \cdot D_{x,3}D_{y,3}D_{z,3}D_{w,3} \cdot x^3zw^3 \\ & + y^2 \cdot D_{x,3}D_{y,3}D_{z,3}D_{w,3} \cdot yz^3w^3 + x^2 \cdot D_{x,3}D_{y,3}D_{z,3}D_{w,3} \cdot xy^3z^3. \end{aligned}$$

8. THE CODE OF THE ALGORITHM

The aim of this final section is to show our implementation in Macaulay2 of the algorithm described in Section 3 of this manuscript; the whole code can be found in [BDSV14]. Throughout this section, $R = \mathbb{F}_p[x_1, \dots, x_d]$ will be the polynomial ring with d variables with coefficients in the field \mathbb{F}_p .

First of all, we write down the code of a procedure which, given an ideal I of R , return as output $I^{[p^e]}$, i.e., the ideal generated by all the p^e th powers of elements in I . The method below is based on code written by M. Katzman and included, among other places, in [KS12b].

```
frobeniusPower(Ideal,ZZ) := (I,e) ->(
R:=ring I;
p:=char R;
local u;
local answer;
G:=first entries gens I;
if (#G==0) then
{
    answer=ideal(0_R);
}
else
{
    N:=p^e;
    answer=ideal(apply(G, u->u^N));
};
answer
);
```

Now, we exhibit the code of a function which, given ideals A, B of R , produces as output the ideal $I_e(A) + B$. For our purposes in this manuscript, $B = (0)$ and A is a principal ideal. Once again, this is based on code written by Katzman and included in [KS12b].

```
ethRoot(Ideal,Ideal,ZZ):= (A,B,e) ->(
R:=ring(A);
pp:=char(R);
F:=coefficientRing(R);
n:=rank source vars(R);
vv:=first entries vars(R);
R1:=F[vv, Y_1..Y_n, MonomialOrder=>ProductOrder{n,n},MonomialSize=>16];
q:=pp^e;
J0:=apply(1..n, i->Y_i-substitute(vv#(i-1)^q,R1));
S:=toList apply(1..n, i->Y_i=>substitute(vv#(i-1),R1));
G:=first entries compress( (gens substitute(A,R1))%gens(ideal(J0)) );
L:=ideal 0_R1;
apply(G, t->
{
    L=L+ideal((coefficients(t,Variables=>vv))#1);
});
L1:=L+substitute(B,R1);
L2:=mingens L1;
L3:=first entries L2;
L4:=apply(L3, t->substitute(t,S));
use(R);
```

```

substitute(ideal L4,R)
);

```

Next, we provide the code of our implementation of Algorithm 3.1. Namely, given $f \in R$, the procedure below gives as output the pair $(e, I_e(f^{p^e-1}))$, where e is the level of f , and $I_e(f^{p^e-1})$ is the ideal where the chain (2.1.1) stabilizes. As the reader can easily point out, this method is just turning Theorem 2.5 into an algorithm.

```

differentialOperatorLevel(RingElement):=(f) ->(
R:=ring(f);
p:=char(R);
local J;
local I;
local e;
e=0;
flag:=true;
local q;
local N;
while (flag) do
{
e=e+1;
q=p^e;
N=q-1;
I=ethRoot(ideal(f^N),ideal(0_R),e);
J=frobeniusPower(I,e);
N=q-p;
if ((f^N)% J==0) then flag=false;
};
(e,I)
);

```

Now, let $x = x_i$ (for some $1 \leq i \leq d$), $n \geq 0$, and $f \in R$. The below procedure returns as output

$$\frac{1}{n!} \frac{\partial^n f}{\partial x^n}.$$

It is worth noting that, in some intermediate step of this method, we have to lift our data to characteristic zero in order to avoid problems with the calculation of $1/n!$.

```

DiffOperator(RingElement,ZZ,RingElement):=(el,numb,funct)->
(
  R:=ring(el);
  vv:=first entries vars(R);
  S:=QQ[vv];
  funct1:=substitute(funct,S);
  el:=substitute(el,S);
  for i to numb-1 do
  funct1=diff(el,funct1);
  funct1=1/(numb!)*funct1;
  funct=substitute(funct1,R);
  el=substitute(el,R);
  use R;
  return funct;
);

```

Next, we provide a method which, given a monomial $\mathbf{x}^\alpha = x_1^{a_1} \cdots x_d^{a_d}$ with $0 \leq a_i \leq p-1$ for all i , returns as output the differential operator $\delta_\alpha \in \mathcal{D}_R^{(e)}$ where, for any other monomial $\mathbf{x}^\beta = x_1^{b_1} \cdots x_d^{b_d}$ with $0 \leq b_i \leq p-1$ for all i , δ_α acts in the following way:

$$\delta_\alpha(\mathbf{x}^\beta) = \begin{cases} 1, & \text{if } \alpha = \beta, \\ 0, & \text{otherwise.} \end{cases}$$

As the reader can easily point out, the below method is just turning Claim 3.4 into an algorithm.

```
DeltaOperator(RingElement,ZZ):=(el,pe)->
```

```
(
  R:=ring(el);
  indet:=vars(R);
  for i to numColumns(indet)-1 do
  (
    deg:=degree(indet_(0,i),el);
    listdiff_i=pe-1-deg;
  );
  return listdiff;
);
```

The next two methods are quite technical; however, both are necessary in order to avoid problems during the execution of our main procedure, which we are almost ready to describe.

```
checkcondition=method();
```

```
--- checkcondition finds a random monomial in startingpol//pol with all the
```

```
--- variables in degree <p^e
```

```
checkcondition(ZZ, RingElement, Ideal, ZZ):=(pe, startingpol, J, indexvar) ->
```

```
(
  pol:=(first entries gens J)_indexvar;
  genJ=first entries gens J;
  R:=ring(pol);
  var:=vars R;
  nvars:=numColumns var;
  for i to numgens J-1 do
    sett_i=set first entries monomials(startingpol//genJ_i);
  supportpol= first entries monomials (startingpol//pol);
  contat=0;
  for i to #supportpol-1 do
  (
    flag=true;
    for k to nvars-1 do
      if degree(var_(0,k),supportpol_i)>pe-1 then flag=false;
      if flag then
        (
          rightsupport_contat=supportpol_i;
        contat=contat+1;
        );
      );
    correctmon=rightsupport_(random contat);
```

```

    return correctmon;
);
-----

DifferentialAction=method();

--- DifferentialAction computes differential(element)

DifferentialAction(RingElement, RingElement):=(differential, element)->
(
    R:=ring(element);
    T:=ring(differential);
    var:=vars T;
    nvars:=numColumns var;
    total=0;
    diffmonomials=first entries monomials (differential);
    for i to #diffmonomials-1 do
    (
multiplication=substitute(element, T);
for j to floor(nvars/2)-1 do
    multiplication=multiplication*
        var_(0, floor(nvars/2)+j)^(degree(var_(0, floor(nvars/2)+j), diffmonomials_i));
differentiation=coefficient(diffmonomials_i, differential)*multiplication;
for j to floor(nvars/2)-1 do
    differentiation=DiffOperator(var_(0, nvars-1-j),
degree(var_(0, floor(nvars/2)-1-j), diffmonomials_i), differentiation);
        total=total+differentiation;
    );
return total;
);

```

We conclude showing our implementation of the algorithm described in Section 3, which is the main result of this paper.

```

DifferentialOperator(RingElement):=(f)->
(
    R:=ring(f);
    p:=char(R);
    F:=coefficientRing(R);
    local e;
    local J;
    (e, J)=differentialOperatorLevel(f);
    J=frobeniusPower(J, e);
    variable:=vars(R);
    variable1:=first entries variable;
    genJ:=first entries gens J;
    nvars:=numColumns vars R;
    powerf:=f^(p^e-1);
    T:=F[d_0..d_(nvars-1), variable1]; --creating ring of differentials
    varDelta:=first entries vars T;
    powerfT=substitute(powerf, T);

```

```

    use R;
    condition:=true;
    while condition do
    (
    for i to numgens J-1 do
    (
    listsupport=checkcondition(p^e,powerf,J,i);
listdiff=DeltaOperator(listsupport,p^e);
    delta1=1;
    delta2=1;
    for j to nvars-1 do
    (
        use T;
        delta1=delta1*varDelta_j^(p^e-1);
        expon2=listdiff_j;
        delta2=delta2*varDelta_(nvars+j)^expon2;
    );
    delta_i=delta1*delta2;
newgen_i=DifferentialAction(delta_i,powerfT);
    );
    newgenmatr=matrix(newgen_0);
    for i from 1 to numgens J-1 do
        newgenmatr=newgenmatr|newgen_i;
    newgenmatrR=substitute(newgenmatr,R);
    if J==ideal(newgenmatrR) then
    (
        matalpha=f^(p^e-p)//newgenmatrR;
        matalpha=substitute(matalpha,T);
        for i to numgens J-1 do
            if i==0 then matrixdiff=matrix{{matalpha_(i,0),delta_i}}
            else matrixdiff=matrixdiff||matrix{{matalpha_(i,0),delta_i}};
        totale=0;
        for i to numgens J-1 do
            totale=totale+matalpha_(i,0)*DifferentialAction(delta_i,powerf);
        powerfp=substitute(f^(p^e-p),T);
    return matrixdiff;
    condition=false;
    );
    );
    use R;
);

```

ACKNOWLEDGEMENTS

This project started during the summer school PRAGMATIC 2014. The authors would like to thank Aldo Conca, Srikanth Iyengar, and Anurag Singh, for giving very interesting lectures and for sharing open problems. They also wish to thank Alfio Ragusa and all the organizers of PRAGMATIC 2014 for giving them the opportunity to attend the school. The authors also thank Josep Àlvarez Montaner and, once again, Anurag Singh and Srikanth Iyengar, for many helpful suggestions concerning the material of this manuscript.

REFERENCES

- [ÀMBL05] J. Àlvarez Montaner, M. Blickle, and G. Lyubeznik. Generators of D -modules in positive characteristic. *Math. Res. Lett.*, 12(4):459–473, 2005. [1](#), [2](#), [3](#)
- [AZ04] M. Aigner and G. M. Ziegler. *Proofs from The Book*. Springer-Verlag, Berlin, third edition, 2004. Including illustrations by Karl H. Hofmann. [11](#)
- [BDSV14] A.F. Boix, A. De Stefani, and D. Vanzo. differentialoperator.m2: a Macaulay2 package for computing a differential operator which acts as the Frobenius homomorphism. Available at <http://atlas.mat.ub.edu/personals/aboix/research.html>, 2014. [18](#)
- [Ber72] I. N. Bernštein. Analytic continuation of generalized functions with respect to a parameter. *Funkcional. Anal. i Priložen.*, 6(4):26–40, 1972. [1](#)
- [BMS08] M. Blickle, M. Mustață, and K. E. Smith. Discreteness and rationality of F -thresholds. *Michigan Math. J.*, 57:43–61, 2008. Special volume in honor of Melvin Hochster. [1](#), [2](#)
- [Bøg95] R. Bøgvad. Some results on D -modules on Borel varieties in characteristic $p > 0$. *J. Algebra*, 173(3):638–667, 1995. [1](#)
- [Can15] E. Canton. A note on injectivity of Frobenius on local cohomology of hypersurfaces. Available at <http://arxiv.org/pdf/1502.03184v1>, 2015. [4](#)
- [FST11] O. Fujino, K. Schwede, and S. Takagi. Supplements to non-lc ideal sheaves. In *Higher dimensional algebraic geometry*, RIMS Kôkyûroku Bessatsu, B24, pages 1–46. Res. Inst. Math. Sci. (RIMS), Kyoto, 2011. [4](#)
- [Gro67] A. Grothendieck. Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas IV. *Inst. Hautes Études Sci. Publ. Math.*, (32), 1967. [2](#)
- [GS] D. R. Grayson and M. E. Stillman. Macaulay2, a software system for research in algebraic geometry. Available at <http://www.math.uiuc.edu/Macaulay2/>. [2](#)
- [Hus87] D. Husemoller. *Elliptic curves*, volume 111 of *Grad. Texts in Math.* Springer-Verlag, New York, 1987. With an appendix by Ruth Lawrence. [10](#)
- [ILL⁺07] S. B. Iyengar, G. J. Leuschke, A. Leykin, C. Miller, E. Miller, A. K. Singh, and U. Walther. *Twenty-four hours of local cohomology*, volume 87 of *Grad. Stud. Math.* American Mathematical Society, Providence, RI, 2007. [1](#), [9](#)
- [KS12a] M. Katzman and K. Schwede. An algorithm for computing compatibly Frobenius split subvarieties. *J. Symbolic Comput.*, 47(8):996–1008, 2012. [4](#)
- [KS12b] M. Katzman and K. Schwede. FSplitting, a Macaulay2 package implementing an algorithm for computing compatibly Frobenius split subvarieties. Available at <http://katzman.staff.shef.ac.uk/FSplitting/>, 2012. [18](#)
- [Luc78] E. Lucas. Sur les congruences des nombres eulériens et les coefficients différentiels des fonctions trigonométriques suivant un module premier. *Bull. Soc. Math. France*, 6:49–54, 1878. [4](#)
- [Ser73] J.-P. Serre. *A course in arithmetic*, volume 7 of *Grad. Texts in Math.* Springer-Verlag, New York-Heidelberg, 1973. [8](#)
- [Yek92] A. Yekutieli. An explicit construction of the Grothendieck residue complex. *Astérisque*, (208):127, 1992. With an appendix by Pramathanath Sastry. [1](#), [2](#)

DEPARTMENT OF ECONOMICS AND BUSINESS, UNIVERSITAT POMPEU FABRA, JAUME I BUILDING, RAMON TRIAS FARGAS 25-27, 08005 BARCELONA, SPAIN.

E-mail address: alberto.fernandezb@upf.edu

URL: <http://atlas.mat.ub.edu/personals/aboix/>

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF VIRGINIA, 141 CABELL DRIVE, KERCHOF HALL CHARLOTTESVILLE, VA 22903, USA

E-mail address: ad9fa@viginia.edu

URL: <http://people.virginia.edu/~ad9fa/>

DEPARTIMENTO DI MATEMATICA E INFORMATICA, UNIVERSITÀ DI FIRENZE, VIALE MORGAGNI, 67/A - 50134 FIRENZE, ITALY

E-mail address: davide.vanzo@unifi.it